










| レジメン | アブラキサン [®] 療法 | | | | | | | | | | | | | | | | | 期間 | 28日間 | | | | | | | | | | | | | |
|----------------------------------|------------------------|-------|---|---|---|---|---|---|---|---|---|---|----|----|----|----|----|-----|------|-----|-----|-----|-----|-----|-----|---|-----|-----|----|--|--|--|
| 薬品名 | 標準投与量 | 投与時間 | day | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | ... | 28 | | | |
| 生食50mL | | 血管確保用 |  | | | | | | | | |  | | | | | | | | | | | | | |  | | | | | | |
| アブラキサン [®] 注 /生食50mL | 100mg/m ² | 30分 |  | | | | | | | | |  | | | | | | | | | | | | | |  | | | | | | |
| 生食50mL | | 15分 |  | | | | | | | | |  | | | | | | | | | | | | | |  | | | | | | |
| 詳細 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |